

Finite temperature phase diagram of the classical Kitaev-Heisenberg model

Craig Price¹ and Natalia B. Perkins²

¹*Department of Physics, The Pennsylvania State University,
104 Davey Lab, University Park, Pennsylvania 16802, USA*

²*Department of Physics, University of Wisconsin,
1150 University Ave., Madison, Wisconsin 53706, USA*

(Dated: April 30, 2013)

We investigate the finite-temperature phase diagram of the classical Kitaev-Heisenberg model on the hexagonal lattice. Due to the anisotropy introduced by the Kitaev interaction, the model is magnetically ordered at low temperatures for all values of parameters at which the model has a discrete symmetry. The ordered phase is stabilized entropically by an order by disorder mechanism where thermal fluctuations of classical spins select collinear magnetic orders in which magnetic moments point along one of the cubic directions. We find that there is an intermediate phase between the low-temperature ordered phase and the high-temperature disordered phase. We show that the intermediate phase is a critical Kosterlitz-Thouless phase exhibiting correlations of the order parameter that decay algebraically in space. Using finite size scaling analysis, we determine the boundaries of the critical phase with reasonable accuracy. We show that the Kitaev interaction plays a crucial role in understanding the finite temperature properties of $A_2\text{IrO}_3$ systems.

I. INTRODUCTION

Recently spin-orbit coupling (SOC) effects have become a subject of intense research across many different disciplines in condensed matter physics. These effects are especially pronounced in 4d and 5d transition-metal compounds whose significant atomic SOC is due to their large atomic weight.

The strongly entangled, spin-orbital nature of the localized states are characterized by an effective angular momenta, J_{eff} . The interactions among the components of J_{eff} are determined by the combination of spin and lattice symmetries, which creates a possibility for unconventional anisotropic exchange interactions and, as a result, a realization of various novel properties.¹⁻³ For example, breaking the spin rotation symmetry permits magnetic Hamiltonians that contain terms that are products of different components of spin operators. Such terms, not allowed in the traditionally studied $SU(2)$ symmetric models, introduce a new source of frustration⁴ and might drive the system towards quantum spin liquid states.

A prominent model exemplifying these kinds of highly anisotropic interactions is the Kitaev model on the honeycomb lattice.⁵ The ground state of the Kitaev model is known exactly; it is a spin liquid characterized by anyonic excitations with exotic fractional statistics. Recently, Jackeli and Khaliullin¹ suggested that the Kitaev model could actually be realized within the honeycomb iridates with general formula $A_2\text{IrO}_3$, since the anisotropic part of the interactions among Ir-ions has the same form as the Kitaev coupling. This suggestion has triggered a lot of experimental⁶⁻⁹ and theoretical¹⁰⁻¹⁶ activity in the study of Na_2IrO_3 and Li_2IrO_3 compounds.

In $A_2\text{IrO}_3$ materials, Ir^{4+} ions are in a low spin $5d^5$ configuration with an effective angular momentum of $J_{\text{eff}} = 1/2$ due to strong SOC. The low-energy Hamiltonian describing the interaction between J_{eff} iridium moments is called the Kitaev-Heisenberg (KH) model

which contains both the isotropic antiferromagnetic (AF) Heisenberg and the anisotropic Kitaev interactions.

The Kitaev exchange interaction is generated through the 90° Ir-O-Ir hopping path between J_{eff} Kramers doublets and is non-zero only in the presence of finite Hund's coupling. At the same time, the isotropic AF Heisenberg interaction is suppressed; the only non-vanishing contribution is from the superexchange interaction arising from the direct overlap of the Ir 5d orbitals. The isotropic exchange via the 90° Ir-O-Ir is canceled due to a destructive interference among multiple superexchange paths.

The ground state of the KH model can be a spin liquid despite the presence of the direct isotropic exchange. This is because the Kitaev spin liquid is rather stable with respect to the Heisenberg interaction^{2,16}; it remains the ground state of the KH model for a wide range of the Heisenberg interaction. Even so, the Kitaev spin liquid in honeycomb iridates has not been observed yet; in both Na_2IrO_3 and Li_2IrO_3 , magnetic order has been observed at low temperatures.⁶⁻⁸ Moreover for Na_2IrO_3 , the KH model in its original form does not appear to be sufficient to account for neither the zigzag magnetic order nor the spectrum of magnetic excitations measured in neutron scattering experiments.⁷⁻⁹

To explain the experimental observations in Na_2IrO_3 , three different modifications of the super-exchange model describing the interactions between localized magnetic moments have been proposed.¹⁷⁻¹⁹ In the first approach,¹⁷ it was shown that the zigzag magnetic order may be stabilized within the KH model by including substantial second and third neighbour antiferromagnetic interactions. The second approach¹⁹ extends the KH model to its full parameter space by including additional hopping processes based on the $t_{2g} - e_g$ hopping along the 90° Ir-O-Ir paths. The main difference between these two approaches is that the role of the Kitaev interaction is minor in the first approach while it still plays the dominant role in the second one. The third approach




FIG. 1: Four possible magnetic configurations: (a) FM ordering; (b) two-sublattice, AF Néel order; (c) stripy order; (d) zigzag order. Open and filled circles correspond to up and down spins.

assumes that Na_2IrO_3 is significantly distorted from the ideal structure; here, the trigonal distortion is the dominant interaction while the SOC is subdominant. This third model actually does not contain the Kitaev term at all, but it does retain the zigzag ground state as one of its magnetic ground states.¹⁸ Thus, the importance of the Kitaev term in the low-energy physics of the A_2IrO_3 compounds remains an open question.

In this paper, we continue to examine the role of the Kitaev interaction in the magnetic properties of A_2IrO_3 systems using the KH model defined in both the original² and the extended¹⁹ parameter space. We argue that the Kitaev interaction's main effect is the reduction of the symmetry of the system from the continuous $\text{SU}(2)$ symmetry to a discrete $\mathbb{Z}_6$ symmetry. The latter is crucial for the finite temperature properties of quasi two-dimensional (2D) iridates since by the Mermin-Wagner theorem²⁰ the 2D magnetic systems with continuous symmetry do not exhibit long-range magnetic order at any finite temperature. However, there is no such constraint on spin systems with discrete symmetry. Thus, we argue that the Kitaev term is responsible for the presence of the long range magnetic order at temperatures which are significantly higher than those that one can expect from a weak inter-layer coupling making A_2IrO_3 compounds weakly three-dimensional.

We also show that the finite temperature properties of the KH model are similar to those of the six-state clock model.^{21,22} The KH model undergoes two continuous phase transitions as a function of temperature. This gives rise to three different phases: a low-T ordered phase with a spontaneously broken $\mathbb{Z}_6$ symmetry, an intermediate critical phase, and a high-T disordered phase. Finite




FIG. 2: Classical energy as a function of α : in the FM phase ($E_{cl}^M = 3 - 5\alpha$, dot-dashed green line), in the zigzag phase ($E_{cl}^Z = -3\alpha + 1$, dotted black line), in the Néel phase ($E_{cl}^N = 5\alpha - 3$ solid blue line), and in the stripy phase ($E_{cl}^S = -\alpha - 1$ dashed red line).

size scaling analysis of our Monte Carlo (MC) simulations confirmed that the intermediate phase is a critical Kosterlitz-Thouless (KT) phase with floating exponents and algebraic correlations.

The rest of the paper is organized as follows. We start our discussion with Sec. II which contains a brief summary of known facts about the classical KH model that will be used in the rest of the paper. In Sec. III, we discuss in detail the results of our numerical simulations. We analyze the discreteness of the low-temperature phase, and then we discuss the criticality of the intermediate phase and show how the finite-size scaling analysis allows us to determine the boundary of the critical phase. We show that the phase transitions are driven by the order by disorder mechanism, in which the thermal fluctuations of classical spins select collinear spin configurations where the magnetic moments all point along one of the cubic directions. We also demonstrate that at particular points of the phase diagram, for which the continuous symmetry is preserved, the ordered state is destroyed at any non-zero temperature. The section ends with a discussion of the finite temperature phase diagram of the classical KH model.

In Sec. IV, we analyze the finite temperature properties of the KH model in its extended parameter space. We show that for the parameters relevant to the Na_2IrO_3 and Li_2IrO_3 compounds, the model exhibits a zigzag magnetic order in agreement with experimental findings. In Sec. V, we show that there is a significant difference between the KH model and the Heisenberg model with a cubic anisotropy. While the later model also exhibits two phase transitions, its intermediate phase is not critical; it is nematic like. Using finite-size scaling analysis, we show that in this case the phase transitions are of the three-states Potts and Ising universality classes. Sec. VI contains a summary of the obtained results and conclusions.




FIG. 3: A low-T distribution of the projections of the vector order parameter on the (111) plane. a) Néel order parameter for $\alpha = 0$; b) Néel order parameter for $\alpha = 0.25$; c) stripy order parameter for $\alpha = 0.5$; d) stripy order parameter for $\alpha = 0.75$.

II. THE CLASSICAL KH MODEL

The classical version of the KH model is

$$\mathcal{H} = -J_K \sum_{\langle ij \rangle_\gamma} S_i^\gamma S_j^\gamma + J_H \sum_{\langle ij \rangle} \mathbf{S}_i \mathbf{S}_j. \quad (1)$$

Following the notation of Ref. 2, we denote the effective degrees of freedom of Ir^{4+} ions as S , and take the quantization axes along the cubic axes of the IrO_6 octahedra. $\gamma = x, y, z$ denotes the three bonds of the honeycomb lattice. Two couplings, J_H and J_K , are opposite in sign; the AF isotropic exchange, J_H , and the ferromagnetic (FM) anisotropic exchange, J_K .

The exchange constants corresponding to the Kitaev and the Heisenberg interactions in the KH model (1) can be conveniently described by one parameter, α , such that $J_K = 2\alpha$ and $J_H = 1 - \alpha$. The model then reads as

$$\mathcal{H} = -2\alpha \sum_{\langle ij \rangle_\gamma} S_i^\gamma S_j^\gamma + (1 - \alpha) \sum_{\langle ij \rangle} \mathbf{S}_i \mathbf{S}_j. \quad (2)$$

Let us briefly describe possible magnetically ordered states of model (2). For this we need to introduce four sublattices: A, B, C , and D . For $0 < \alpha < 1/3$, the classical ground state is the simple two-sublattice AF Néel order (Fig.1 (b)) characterized by the order parameter

$$\mathbf{N} = \frac{1}{\mathcal{N}} \sum_i (\mathbf{S}_{iA} - \mathbf{S}_{iB}). \quad (3)$$

$\mathcal{N}$ denotes the number of sites. The stripy AF order (Fig.1 (c)) describes the classical ground state of the model for $1/3 < \alpha < 1$ and is given by

$$\mathbf{S} = \frac{1}{\mathcal{N}} \sum_i (\mathbf{S}_{iA} - \mathbf{S}_{iB} - \mathbf{S}_{iC} + \mathbf{S}_{iD}). \quad (4)$$

For $\alpha = 1.0$, the stripy AF state is classically degenerate with other magnetically ordered states (see Fig.2). For example, it is degenerate with FM order (Fig.1 (a)) described by the total magnetization

$$\mathbf{M} = \frac{1}{\mathcal{N}} \sum_i \mathbf{S}_{iA}, \quad (5)$$

and with zigzag AF spin order (Fig.1 (d)) described by

$$\mathbf{Z} = \frac{1}{\mathcal{N}} \sum_i (\mathbf{S}_{iA} + \mathbf{S}_{iB} - \mathbf{S}_{iC} - \mathbf{S}_{iD}). \quad (6)$$

The classical degeneracy of the point corresponding to the classical Kitaev model is macroscopic and is known exactly; asymptotically, it has $(1.662)^N$ spin configurations. This was computed by Baskaran *et al*²³ by mapping ordered states of the classical Kitaev model to a certain dimer covering of the honeycomb lattice whose total number is known.²⁴

Because of the presence of the anisotropic Kitaev interaction term, the KH model has discrete spin-rotation symmetry for all of α except at two points, $\alpha = 0$ and $\alpha = 0.5$. At $\alpha = 0$, the KH model reduces to the antiferromagnetic Heisenberg model with continuous $\text{SU}(2)$ symmetry. At $\alpha = 0.5$, the stripy phase becomes an exact ground state of the model; it corresponds to a fully polarized FM state in a rotated basis. This basis can be seen by fixing the spin's direction on sublattice A and rotating the spins on sublattices B, C , and D by the angle π about the x, y , and z axis, respectively (see Fig. 1(c)).² It is evident that the FM state has true $\text{SU}(2)$ symmetry. Away from these special points, the symmetry of the KH model is discrete; it combines the cubic symmetry of both the spin and the lattice space.

III. FINITE TEMPERATURE PHASE DIAGRAM AND CRITICAL PROPERTIES OF THE CLASSICAL KH MODEL

In this section we study the behavior of the classical KH model (2) at finite temperature using MC simulations based on the standard Metropolis algorithm. In our simulations, we treat the spins as three-dimensional (3D) vectors, $\mathbf{S}_i = (S_i^x, S_i^y, S_i^z)$, of unit magnitude with $(S_i^x)^2 + (S_i^y)^2 + (S_i^z)^2 = 1$. At each temperature, more than 10^7 MC sweeps were performed. Of these, 10^6 were used to equilibrate the system, and afterwards only 1 out of every 5 sweeps was used to calculate the averages of physical quantities. We present all energies in units of J_H and assume $k_B = 1$. The simulations were performed




FIG. 4: Snapshots of spin configurations in the low-T phase computed for a) $\alpha = 0$, b) $\alpha = 0.25$ c) $\alpha = 0.5$ and d) $\alpha = 0.75$. Each color corresponds to one of the two or four different sublattices describing the Néel or the stripy order parameters, respectively.

on different systems with a total number of sites equal to $2 * L * L$. The systems are spanned by the primitive vectors of a triangular lattice $\mathbf{a}_1 = (1/2, \sqrt{3}/2)$ and $\mathbf{a}_2 = (1, 0)$ with a 2-point basis using periodic boundary conditions.

In our simulations we compute the following observables: four different order parameters $\mathbf{O} = \{\mathbf{N}, \mathbf{S}, \mathbf{M}, \mathbf{Z}\}$ defined in Eqs.(3-6), corresponding susceptibilities

$$\chi_{\mathbf{O}} = \mathcal{N}(\langle \mathbf{O}^2 \rangle - \langle \mathbf{O} \rangle^2)/T, \quad (7)$$

the Binder's cumulants

$$B_{\mathbf{O}} = 1 - \langle \mathbf{O}^4 \rangle / 3 \langle \mathbf{O}^2 \rangle^2, \quad (8)$$

and the specific heat

$$C = (\langle E^2 \rangle - \langle E \rangle^2) / \mathcal{N} T^2. \quad (9)$$

A. Low-temperature ordered phase

At low temperatures, the KH model magnetically orders in either a Néel state, $\mathbf{N}$, or in a stripy state, $\mathbf{S}$, depending on the relative strengths of the Kitaev and the Heisenberg interactions. The presence of long range order at finite temperatures requires a discreteness of the order parameter, and means that the direction of the order parameter must also be selected. This, however, does not happen on the level of non-interacting spin waves. For both the Néel state and the stripy state, the linear spin-wave spectrum has a quasi-Goldstone mode at the ordering vector simulating the spontaneous breaking of the continuous symmetry.² The discrete symmetry of the order parameter appears only due to the contribution of higher order anharmonic modes of spin fluctuations which remove the accidental degeneracy of the classical ground states and lower the energy of the states whose order parameter points along cubic directions.² This gives

a six-fold degeneracy of the order parameter manifold which corresponds to the six degenerate minima in the free energy. This is a well known order by disorder mechanism in which spin fluctuations (quantum or thermal) remove the accidental degeneracy and select the true ordered state.

The discreteness of the order parameter at all but special points can be revealed from the histogram method in which we record two-dimensional distributions of the projections of the vector order parameter on the (111) plane. We use the projection of the order parameter on the (111) plane because it preserves the cubic symmetry of the model and shows that its degeneracy is related to the orientation of the order parameter with respect to the directions in a cubic crystal. In Fig. 3 (a), (b), (c), and (d) we present, respectively, the low-T distributions of the projection of the vector order parameter on the (111) plane for four different values of α : 0, 0.25, 0.5, and 0.75. The six-fold peak structure is observed in the distribution function of the Néel and the stripy order parameters for $\alpha = 0.25$ (Fig. 3 (b)) and $\alpha = 0.75$ (Fig. 3 (d)). The peaks correspond to the ordered spin configurations with an order parameter pointing along one of the cubic axes. In Figs. 4 (b) and (d) we present snapshots of the spin configurations which contribute to the histograms in Figs. 3 (b) and (d). In the snapshot of the spin configuration shown in Fig. 4(b), the spins that are pointing along the y -direction are antiparallel on the two sublattices. This state corresponds to the Néel phase with the order parameter directed along y -axis. In the spin configuration of Fig. 4 (d), spins are pointing parallel to the z -direction on two sublattices and antiparallel on another two. This state corresponds to the stripy phase with the order parameter directed along z -axis.

The projections and snapshots of the order parameter for the two special points $\alpha = 0$ and $\alpha = 0.5$ are presented in Figs. 3 (a) and (c) and Figs. 4 (a) and (c),




FIG. 5: Histograms of the order parameter $m_{N(S)}$, obtained for the system with $2*84*84$ spins in the ordered phase, (a) and (e), in the intermediate phase, (b)-(c) and (f)-(g), and in the disordered phase, (d) and (h). Histograms (a)-(d) are computed for $\alpha = 0.25$, and (e)-(h) are for $\alpha = 0.75$. The histograms are presented on the complex plane ($\text{Re} |m_{N(S)}|$, $\text{Im} |m_{N(S)}|$).

respectively. These are the points with continuous symmetry. As expected, we see that the projections of the order parameter on the (111) plane (3 (a) and (c)) are more or less equally distributed along a circle. Note that the multiple "quasi"-peaks appear because the computations are performed on a finite size system. The snapshot for $\alpha = 0$ (4 (a)) shows that a certain direction is chosen in this particular spin configuration but it is not along one of the cubic axes. Because this direction varies from configuration to configuration and also because the spins belonging to the same sublattice can be orientated parallel or antiparallel to this direction, there is no long range order for $\alpha = 0.0$. A similar situation is observed at $\alpha = 0.5$. Here, spins at each sublattice point along a certain direction which is not chosen to be along any cubic axes. There is no long range magnetic at $\alpha = 0.5$ as well.

Finally, there is no additional degeneracy of the order parameter related to its real-space structure. Both the stripy and the zigzag order can have three different configurations of bond orderings due to the 120° rotational symmetry of the honeycomb lattice. The absence of the additional degeneracy can be understood from the fact that, due to SOC, the symmetry transformations act simultaneously on spins and on the lattice. Once the pattern of the bond ordering is chosen, the direction of the order parameter in spin space is also chosen. For example, in the vertical stripy state shown in Fig. 1(c), the spins are directed along the z axis just like in the snapshot shown in Fig. 4 (c).

B. The critical nature of the intermediate phase. Finite size scaling analysis.

Regardless of the specific kind of magnetic order, the low-T ordered phase is separated from the high-T paramagnetic phase by the intermediate phase. In our recent study, we have shown that the intermediate phase is a critical phase with two finite-temperature boundaries that correspond to Berezinskii-Kosterlitz-Thouless (BKT) phase transitions.¹⁴

To describe the finite temperature properties of the KH model, we use a projection of the vector order parameter on the (111) plane. The vector is characterized by both the absolute value and the azimuthal angle of $|m_{N(S)}|$. This projection is equivalent to the $\mathbb{Z}_6$ order parameter, and we write it in a complex form via $m_{N(S)} = \sum_{i=1}^6 |m_{i,N(S)}| e^{i\theta_i}$. We chose the phase θ such that the minimal-energy states of the order parameter, which point along the cubic axes, will be labeled by the values $\theta_i = \pi n_i/3$, $n_i = 0, \dots, 5$. Observing the critical phase proved to be challenging for several reasons. First, within the vicinity of the BKT transition, the critical behavior gives rise to a very slow, logarithmic convergence to the thermodynamic limit. We thus had to perform the finite size scaling analysis of our simulations on rather large systems with $L = 84, 96, 108, 120, 144, 168, 204$. Second, contrary to the Ising-like spin systems in which all previous studies of the six-state clock model's critical phase have been performed, the magnetic degrees of freedom in the classical KH model are 3D Heisenberg spins which are affected by thermal fluctuations more strongly. As a result, a large number of sweeps is needed to aver-




FIG. 6: The log-log plots of the order parameter $m_{N(S)}$ as a function of system size L at various temperatures. The solid curves indicate the linear behavior that corresponds to a power-law dependence, $m_{N(S)} \sim L^{-\eta/2}$, corresponding to the intermediate critical phase. The dashed curves show deviation away from the linear behavior outside the critical phase.




FIG. 7: The Binder cumulant as a function of temperature for (a) $\alpha = 0.25$ and (b) $\alpha = 0.75$. From the crossing points of different Binder's curves, we estimate $T_{c1} = 0.152$ and $T_{c1} = 0.124$ for $\alpha = 0.25$ and $\alpha = 0.75$, respectively.

age out the thermal fluctuations and capture the critical behavior.

A clear indication of the three-phase structure emerges from the inspection of the histogram plot of the complex magnetization $m_{N(S)}$ at different temperatures (see, Fig.5). As we move from low to high temperature, we observe a transition from an ordered phase (6 isolated spots) through an intermediate critical phase (ring distribution) up to the disordered phase (uniform distribution around zero). The critical phase has an emergent, continuous $U(1)$ symmetry,²⁵ which is reminiscent of the intermediate phase of the six-state clock model.²¹ At the

boundaries, T_{c1} and T_{c2} , and inside the critical phase, the order parameter exhibits a power law dependence on system size of the form, $m_{N(S)} \sim L^{-\eta/2}$. From renormalization group analysis,²¹ it is known that the lower T_{c1} and the upper T_{c2} transitions in the six state clock model are characterized by the critical exponents $\eta_1 = 1/9$ and $\eta_2 = 1/4$, respectively.

Fig. 6 shows the log-log plots of the order parameter m_N and m_S as a function of system size L for different temperatures. In accordance with the power law behavior of the order parameter, the data points of the log-log plot show a linear behavior inside the critical phase. For $\alpha = 0.25$ (Fig. 6 (a)), the log-log plots of the order parameter as function of L show a linear behavior in the temperature interval between $T_{c1} \simeq 0.152$ and $T_{c2} \simeq 0.162$, with the boundaries characterized by the critical exponents $\eta_1 = 0.13$ and $\eta_2 = 0.21$. For $\alpha = 0.75$, we detected the critical phase in the temperature interval between $T_{c1} \simeq 0.125$ ($\eta_1 = 0.13$) and $T_{c2} \simeq 0.127$ ($\eta_2 = 0.22$). For both $\alpha = 0.25$ and $\alpha = 0.75$, our estimates for the critical exponents of the boundaries are in a good agreement with theoretical values for the six state clock model.

We can verify our estimates for the BKT transition temperatures using the Binder cumulants method and further finite-size scaling analysis.³¹ The Binder cumulants $B_{m_{N(S)}}$, defined in Eq.(8), have a scaling dimension of zero; thus the crossing point of the cumulants for different lattice sizes provides a reliable estimate for the value of the critical temperature T_{c1} at which the long range order is destroyed. The results for the Binder cumulants $B_{m_{N(S)}}$ are presented in Fig. 7. The crossing points for $\alpha = 0.25$ and $\alpha = 0.75$ are at $T_{c1} = 0.153$ and at $T_{c1} = 0.125$, respectively, and are in a good agreement with estimates obtained from the log-log plots in Fig. 6.

Another way to check the nature of the critical phase boundaries is by using the data collapse method based on the finite size scaling arguments. In the BKT transition, the order parameter and its susceptibility exhibit power law behavior, $m_{N(S)} \sim \xi^{-\eta/2}$ and $\chi_{N(S)} \sim \xi^{2-\eta}$, while the correlation length near the critical temperature, T_c , diverges as $\xi \sim \exp(at^{1/2})$, where a is a non-universal constant and $t = |T - T_c|/T_c$ is the reduced temperature.³² The finite size scaling analysis is based on the assumption that the singular part of the free energy is a homogeneous function of system size L and correlation length ξ , and depends only on their ratio L/ξ . Based on this assumption, the finite size scaling behavior of the order parameter and the susceptibility have the following functional forms

$$m_{N(S)} = L^{-b} M_{N(S)} \left(\frac{L}{\xi} \right) \quad (10)$$

$$\chi_{m_{N(S)}} = L^c \Xi_{m_{N(S)}} \left(\frac{L}{\xi} \right),$$

where scaling constants $b = \eta/2$ and $c = 2 - \eta$, and $M_{N(S)}$ and $\Xi_{m_{N(S)}}$ are unknown universal functions. We plot the variation in temperature of the order parameter




FIG. 8: Thermodynamic quantities and finite size scaling analysis. (a) and (b): The order parameters m_N and m_S as functions of T for $\alpha = 0.25$ and $\alpha = 0.75$, respectively. (c) and (d): The susceptibility, $\chi_{m_{N(S)}}$ (defined in Eq.(7)) plotted as a function of T for $\alpha = 0.25$ and $\alpha = 0.75$, respectively. (e) and (f): Finite-size scaling of the order parameter data in the low-temperature region ($T < T_{c1}$) for $\alpha = 0.25$ and $\alpha = 0.75$, respectively. (h) and (g): Finite-size scaling of the high-temperature susceptibility data ($T > T_{c2}$) for $\alpha = 0.25$ and $\alpha = 0.75$, respectively.

in Fig.8 (a) and (b) and the susceptibility vs temperature in Fig.8 (c) and (d). In Fig.8 (e) and (f) and Fig.8 (h) and (g), we show the scaling plots for $m_{N(S)}L^b$ as a function of $L^{-1} \exp(a/\sqrt{(T_{c1}-T)/T_{c1}})$ and $\chi_{m_{N(S)}}L^{-c}$ as a function of $L^{-1} \exp(a/\sqrt{(T-T_{c2})/T_{c2}})$, respectively.

The finite-size effects in the behavior of the order parameter and its susceptibility are striking: we see both a significant finite-size tail of $m_{N(S)}$ into intermediate region for $T > T_{c1}$ and a strong dependence of the position and height of the susceptibility peak on the system size L . This is a consequence of the infinite correlation length in the intermediate critical region, which in the finite size systems look like the quasi long range order. Nevertheless, the data points for different system sizes plotted in the scaled form collapse reasonably well on universal curves that correspond to the universal functions $M_{N(S)}$ and $\Xi_{m_{N(S)}}$. The best data collapse was obtained for the following scaling parameters: $a = 1.9$, $b = 0.056$ and $c = 1.45$ and transition temperatures $T_{c1} = 0.153$ and $T_{c2} = 0.1615$ for $\alpha = 0.25$; $a = 1.55$, $b = 0.056$ and $c = 1.55$ and transition temperatures $T_{c1} = 0.125$ and $T_{c2} = 0.127$ for $\alpha = 0.75$. The values obtained for b and c give the following critical exponents for the lower and the upper boundary of the critical phase: $\eta(T_{c1}) = 0.11$, $\eta(T_{c2}) = 0.275$, and $\eta(T_{c1}) = 0.11$, $\eta(T_{c2}) = 0.225$ for $\alpha = 0.25$ and $\alpha = 0.75$, respectively.

To summarize the discussion of the numerical data presented above, we can say that we definitely observe the critical intermediate phase in the classical KH model, which, however, slightly deviates from the standard BKT criticality. We believe that the imperfect data collapse to

be caused by the presence of three dimensional spin fluctuations of Heisenberg spins and finite sized system sizes. Nevertheless, the smallness of the discrepancy and overall high quality of the scaling indicate that this effect is subdominant in the temperature range where the critical behaviour occurs.

C. The finite temperature phase diagram

In Fig.9 we present the finite temperature phase diagram of the KH model (2) obtained by tracking the dependence of the transition temperatures T_{c1} and T_{c2} on the strength of the KH model parameter α . As discussed in detail in the previous section, in order to determine critical temperatures we have used the Binder's cumulant method and finite size scaling arguments.

As expected, only two low-T magnetically ordered phases are present on the phase diagram - the Néel phase at $\alpha < 1/3$ and the stripy AFM phase at $1/3 < \alpha < 1$. The vertical line that separates the Néel and the stripy AFM phases corresponds to a first-order phase transition. Our numerical results show that the critical phase is present for all values of α except the special points $\alpha = 0, 0.5$, and 1 at which the model has continuous symmetry, and the ordered state is destroyed at any non-zero temperature in accordance with Mermin-Wagner theorem.²⁰ Note that the way the width of the critical phase disappears in the vicinity of $\alpha = 0$ and $\alpha = 0.5$ is different than for $\alpha = 1$. In Fig.9, we can see that the critical phase narrows very rapidly in the vicinity of




FIG. 9: Phase diagram of the classical KH model (2). The regions designated by, “N”, and “S”, are the regions of the phase diagram where the Néel, and stripy order persist. The vertical line between the Néel and the stripy phase at $\alpha = 1/3$ denotes the first order phase transition. Each red circle designates a lower critical temperature T_{c1} for the value of α for which the model was explicitly simulated; the red line extrapolates between these points. For each α , the critical value of T_{c1} was determined through the crossings of Binder’s cumulant curves. The critical phase is shown in blue. The upper boundary of the blue critical region was determined by finding the temperature T_{c2} , for which the value of the critical exponent η exceeds 0.25. The blue line extrapolates between each computed point.

$\alpha = 0, 0.5$. This behavior is related to the fact that at these particular points the anisotropy either does not exist as at $\alpha = 0$ or cancels out as at $\alpha = 0.5$. In the vicinity of these points, the local minima of the free energy that corresponds to different directions of the order parameter are still separated by finite energy barriers. These barriers allow finite temperature ordering through the order by disorder mechanism. At the same time, the width of the critical phase slowly decreases when we approach the Kitaev limit and is extremely small in the vicinity of $\alpha = 1$. In this limit, the difference between the classical energies of the stripy, zigzag and ferromagnetic phases is decreasing with the critical temperature tending to zero. At $\alpha = 1$, all of the phases are degenerate (see also Fig. 2 and the discussion above), thus, there is no order-by-disorder in this limit.

IV. FINITE TEMPERATURE PHASE DIAGRAM OF THE EXTENDED CLASSICAL KH MODEL

Here we explore the finite temperature phase diagram of the KH model extended to its full parameter space. This extension, consisting in taking into account all super-exchange processes leading to the coupling between Ir ions, gives a possibility to capture the zigzag magnetic order observed in the $A_2\text{IrO}_3$ compounds.^{7,8}

There are three physically distinct processes that determine the ratio between the Kitaev and the isotropic

Heisenberg terms in the original KH model. Two of the processes involve the virtual hopping of the t_{2g} electrons through the two nearest oxygen ions. As it was shown by Jackeli and Khaliulin², the processes via the upper and lower oxygen interfere destructively and the isotropic part of the Hamiltonian exactly vanishes. Exchange couplings of neighboring Kramers states on Iridium ions appear due to either the multiplet structure of the excited levels on a Ir ion induced by Hund’s coupling, or due to Coulomb interactions. The third process involves a direct hopping between NN t_{2g} orbitals which gives a finite Heisenberg term in the KH model. All of these processes involve only t_{2g} electrons. However, there is another possible process:¹⁹ the intersite $t_{2g} \leftrightarrow e_g$ hopping along the 90° Ir-O-Ir paths. This is the dominant pathway in a 90° geometry since it involves strong $t_{pd\sigma}$ overlap between the p -orbitals on oxygen and the e_g orbitals on iridium ions. Remarkably, these hopping processes also introduce the Kitaev interaction, but with a different sign.

Following Ref.19, we consider the most general extension of the KH model, in which both Kitaev and Heisenberg interactions can change sign and be both FM and AFM. The Hamiltonian of the extended model is

$$\mathcal{H} = A(2 \sin \phi \sum_{\langle ij \rangle_\gamma} S_i^\gamma S_j^\gamma + \cos \phi \sum_{\langle ij \rangle} \mathbf{S}_i \mathbf{S}_j), \quad (11)$$

where the relative strengths of the effective interaction between Ir magnetic moments are described by the phase angle, ϕ defined in the interval $(0, 2\pi)$, and the overall energy scale, $A = \sqrt{J_K^2 + J_H^2}$. The phase space of the original KH model is covered by the values of ϕ from $3\pi/2$ to 2π .

Using MC simulations, we investigate the finite temperature properties of this extended model (11), and present the finite temperature phase diagram in Fig.10. In its full parameter space, the extended KH model accommodates four different classical phases. In addition to the Néel and the stripy phases present in the phase diagram (Fig.9), the FM and the zigzag phases appear in Fig.10 in a wide range of values of ϕ . The previously missing zigzag magnetic order is found to occupy almost a quarter of the phase space of the extended model. Our findings are in agreement with the ground state phase diagram of the quantum model (11) obtained by exact diagonalization.¹⁹ However, as we already discussed above, the classical model does not support the spin liquid phases present in the quantum model.

The finite temperature properties of the extended model are very similar to the finite temperature properties of the original KH model. Both the transition between the zigzag and the FM order, and the transition between the Néel and the stripy are first order phase transitions. The intermediate phase is present for the whole parameter space of the extended model except for special points at which the continuous symmetry is restored. The similarities between finite temperature properties of the original and the extended KH model are not surprising. The above mentioned 4-sublattice spin




FIG. 10: Phase diagram of the extended classical KH model (11). The regions designated by, “N”, “S”, “Z”, and “F” are the regions of the phase diagram where the Néel, stripy, zigzag, and FM order persist. Each red circle designates a lower critical temperature T_{c1} for the value of ϕ for which the model was explicitly simulated. For each value of ϕ , the critical value of T_{c1} was determined through the crossings of Binder’s cumulant curves; the red line extrapolates between these points. The critical phase is shown in blue. The upper boundary of the blue critical region was determined by finding the temperature T_{c2} , for which the value of the critical exponent η exceeds 0.25. The blue line extrapolates between each computed point.

transformation,² permits a mapping of the AF stripy phase to the FM phase and of the AF Néel to the zigzag phase. This implies that these phases have identical dynamics at both zero temperature¹⁹ and at finite temperatures, which is in agreement with our numerical findings.

According to the fit of the uniform magnetic susceptibility presented in Ref.¹⁹, $\phi \simeq 111^\circ \pm 2^\circ \simeq 0.62\pi \pm 0.01\pi$ for Na_2IrO_3 and $\phi \simeq 124^\circ \pm 4^\circ \simeq 0.69\pi \pm 0.03\pi$ for Li_2IrO_3 . Using these values, we get the following estimates for the Néel temperatures: $T_{c1} \simeq 0.16$ which is equivalent to about 17.7 K for Na_2IrO_3 , and $T_{c1} \simeq 0.19$ which is equivalent to about 21 K for Li_2IrO_3 . Both values are close to the experimental value $T_N \simeq 15$ K obtained for both Na_2IrO_3 and Li_2IrO_3 compounds.^{6,8} Our estimates for the upper boundary of the critical phase is $T_{c2} = 0.18$ for Na_2IrO_3 and $T_{c2} = 0.22$ for Li_2IrO_3 which is equivalent to about 20 K and 24.5 K, respectively. This gives the estimates for the width of the critical phase about 2.3 K and 3.5 K for Na_2IrO_3 and Li_2IrO_3 , respectively.

V. CLASSICAL HONEYCOMB HEISENBERG ANTIFERROMAGNET WITH A CUBIC ANISOTROPY

To complete this study, we also investigate the nature of the finite-temperature phase transitions of the honeycomb Heisenberg antiferromagnet in which the cubic anisotropy is included explicitly. In this case, the model

is defined by the Hamiltonian

$$\mathcal{H}_c = J_H \sum_{\langle ij \rangle} \mathbf{S}_i \mathbf{S}_j - D \sum_i ((S_i^x)^4 + (S_i^y)^4 + (S_i^z)^4) \quad (12)$$

where D denotes the strength of the cubic anisotropy. The positive sign of the anisotropy, $D > 0$, determines that spins tend to align along the cubic axes, and not in the diagonal directions of the lattice which would correspond to $D < 0$. Because of the cubic anisotropy, the model (12) shows no full rotational symmetry. The spin projections in the (111) plane again will favor six directions as in the case of the KH model. Here we explore by MC simulations whether or not the finite temperature properties of models (1) and (12) are similar.

Let us discuss the thermodynamic properties of the model (12). At $T = 0$ and in the absence of the anisotropy, the ground state of the model (12) is determined by the Néel order parameter $\mathbf{N}$. In the presence of the anisotropy D , the ordered state will survive until some critical temperature. At high temperature, the thermal fluctuations of spins will destroy the magnetic order and the system will be in the disordered paramagnetic state. As in the KH model, in order to reach the ordered state from high temperature, the system has to break the discrete $\mathbb{Z}_6$ symmetry of the Heisenberg honeycomb model with cubic anisotropy. In the case of the KH model, the $\mathbb{Z}_6$ symmetry was broken through the critical intermediate phase. In our simulations of (12), we did not observe any critical phase; instead, the $\mathbb{Z}_6$ symmetry is broken in the following two steps. After lowering the temperature, the $\mathbb{Z}_3$ symmetry is broken first at $T = T_{c2}$: the system still remains paramagnetic since the expectation value of $\langle \mathbf{N} \rangle = 0$. In the second step, the remaining $\mathbb{Z}_2$ symmetry is broken at the temperature $T = T_{c1} < T_{c2}$ and the system acquires long-range magnetic order.

The numerical results computed for the value of the cubic anisotropy $D = 0.1$ are presented in Figs. 11 (a)-(d). They display the data for the temperature dependence of the staggered magnetization $\mathbf{N}$ and the cubic parameter b_N and corresponding susceptibilities. In our simulations, the cubic order parameter b_N is expressed as the expectation value of a doublet given by

$$b_{N1} = (N_x^2 + N_y^2 - 2N_z^2)/\sqrt{6}, \quad (13)$$

$$b_{N2} = (N_x^2 - N_y^2)/\sqrt{2}, \quad (14)$$

where N_x, N_y, N_z are the components of the Néel order parameter $\mathbf{N}$. At high temperature, the cubic symmetry is not broken and $N_x^2 = N_y^2 = N_z^2$ and $\langle b_{N1} \rangle = \langle b_{N2} \rangle = 0$. As we can see from the Figs. 11, both the order parameters and their susceptibilities indicate a continuous phase transition at different temperatures T_{c1} and T_{c2} for $\mathbf{N}$ and b_N , respectively. At the higher temperature, T_{c2} , the cubic symmetry is spontaneously broken and one of the cubic axes is selected, say z . Then $\langle b_{N1} \rangle = -\sqrt{2/3}$, $\langle b_{N2} \rangle \simeq 0$ and the cubic parameter $\langle \mathbf{b}_N \rangle$ acquires a finite value. Nevertheless, there is still no long range order




FIG. 11: The magnetization N (a) and the cubic order parameter b_N (b) as functions of T . The susceptibilities χ_N (c) and χ_{b_N} (d) as functions of T . (e) and (f): Finite-size scaling of the order parameters N and b_N in the low-temperature region $T < T_{c1}$. (h) and (g): Finite-size scaling of the high-temperature susceptibilities χ_N and χ_{b_N} in the high-temperature region $T > T_{c2}$. The following values of the parameters are used: The anisotropy constant is considered to be $D = 0.1$.

as the time reversal symmetry remains unbroken. Thus, we can say that the intermediate phase is nematic-like. At the lower temperature, T_{c1} , the time reversal symmetry is also broken and the system acquires long-range magnetic order characterized by non-zero $\langle \mathbf{N} \rangle$. We can estimate T_{c1} and T_{c2} using the associated Binder's cumulants, B_N and B_{b_N} , whose temperature dependencies are presented in Figs.12 (a)-(b). From their crossing points we estimated the transition temperatures to be equal to $T_{c1} = 0.314$ and $T_{c2} = 0.32$.

Next, in order to obtain the critical exponents characterizing these two phase transitions we perform a finite-size scaling analysis. As the $\mathbb{Z}_6$ symmetry of the model (12) is reduced to $\mathbb{Z}_2$ symmetry in the intermediate phase, the high- T transition is expected to be in the universality class of 3-states Potts model and the low- T transition to belong to the Ising universality class.

First, using the scaling relation for the Binder cumulants, B_N and B_{b_N} , we obtained the correlation length exponents ν_1 and ν_2 describing the divergence of the correlation length, $\xi \sim |T - T_c|^{-\nu(T_c)}$, close to the critical points $T_c = T_{c1}$ and $T_c = T_{c2}$, respectively. In accordance with the theory prediction, the best data collapse is obtained for $\nu_1 = 0.83$ and $\nu_2 = 1.0$ which are the critical exponents of the three-states Potts and 2D Ising model, respectively.

Second, having found the critical exponents ν_1 and ν_2 , we can estimate the critical exponents β and γ by performing a scaling fit of the order parameters and susceptibilities (Figs. 11 (e)-(g)). Near the critical temperatures, T_{c1} and T_{c2} , the cubic order parameter and staggered magnetization are expected to satisfy the scaling

relation $b_N \simeq L^{-\beta_1/\nu_1}$ and $N \simeq L^{-\beta_2/\nu_2}$, respectively. The scaling laws for their susceptibilities are given by $\chi_{b_N} \simeq L^{\gamma_1/\nu_1}$ and $\chi_N \simeq L^{\gamma_2/\nu_2}$. The best scaling is obtained for the exponents $\beta_1 = 1/9$ and $\gamma_1 = 13/9$ – the exact scaling coefficients for the Potts's transition, and $\beta_2 = 1/8$ and $\gamma_2 = 7/4$ – the exact scaling coefficients for the Ising transition.

To conclude this section, in Fig.13 we present a finite temperature phase diagram for the model (12). We see that the width of the intermediate phase decreases with the increase of the value of the cubic anisotropy parameter D . At around $D \simeq 0.5$, two transitions become indistinguishable and the intermediate phase collapses. The transition between the low- T ordered magnetic phase and high- T disordered phase becomes first order.

VI. CONCLUSION

In this paper we studied finite temperature properties of the classical two-dimensional KH model and computed the phase diagram of this model in its full parameter space. We started by analyzing the lowest-energy magnetic configuration and then found that all of the magnetic phases are accompanied by an accidental continuous rotational degeneracy which does not correspond to any symmetry of the Hamiltonian. This pseudo degeneracy is lifted by thermal fluctuations of spins giving rise to an ordering at low temperature as observed by our MC simulations. Specifically we determined that the low temperature phase is magnetically ordered at all values of parameters at which the model has discrete symme-




FIG. 12: (a) and (b): The Binder cumulants, $B_N = 1 - \langle N^4 \rangle / 3 \langle N^2 \rangle^2$ and $B_{b_N} = 1 - \langle b_N^4 \rangle / 3 \langle b_N^2 \rangle^2$, as functions of T . (c) and (d): Finite-size scaling of the Binder cumulants B_N and B_{b_N} . The anisotropy constant is considered to be $D = 0.1$.




FIG. 13: Finite temperature phase diagram for the classical honeycomb Heisenberg antiferromagnet with the cubic anisotropy (12). The blue line denotes the high-T three state Pott's transition, and the red line denotes the low-T Ising transition.

try. The direction of the order parameter is chosen by thermal fluctuations of the spins through the order by disorder mechanism.

From numerical MC simulations, we have verified that the classical KH model effectively behaves like the six-state clock model. At finite temperatures, the model exhibits two phase transitions and an intermediate phase

between them. Based on the finite size scaling analysis, we have shown that the intermediate phase is the critical phase with algebraically decaying correlations of the order parameter. We found that the phase boundaries of the critical phase are of the BKT type. We also obtained that the numerical values of the critical exponent, η , characterizing these two transitions are compatible with the theoretical expectations based on the renormalization group analysis for the six-state clock model.²¹ It should be emphasized that the mapping of the KH model onto the six state clock model is valid only below a certain temperature which determines the 2D-3D crossover. The crossover temperature is, however, larger than the temperature of the second BKT transition and, therefore, the low-temperature properties of the KH model are captured by the six-state clock model. At high temperatures, when T is larger than any energy scale in the system, the effect of thermal fluctuations is to destroy any kind of order and to put the KH model into the 3D paramagnetic state. In this regime the mapping of the classical KH model to the six state-clock model is not valid.

We also performed a comparative study of honeycomb Heisenberg antiferromagnet with the cubic anisotropy (12). In this case, the continuous symmetry is explicitly broken, which allows for the magnetically ordered phase to persist up to a finite temperature. We have shown that this model has distinct finite temperature properties and that the low-T ordered phase is destroyed in a different way. The similarity between the honeycomb Heisenberg antiferromagnet with the cubic anisotropy and the KH model is that in both cases the ordered phase is destroyed in two steps. The main differences between the two is that when the cubic anisotropy is taken into account explicitly, the intermediate phase is nematic-like, and the phase transitions are in the three-states Potts and 2D Ising universality classes.

Finally, we consider the implications of our numerical results to Na_2IrO_3 and Li_2IrO_3 compounds. We have shown that finite temperature magnetic properties of these systems can be captured by the extended KH model. We found that for the values of the model parameters relevant to these systems, the model exhibits zigzag magnetic order. Also, our numerical estimates of the Néel temperature are close to the experimental values.

Acknowledgements. The authors are particularly thankful to C. Batista, G.-W. Chern, G. Jackeli, Y. Kato and Peter Wölfle for stimulating discussions and many helpful suggestions. N.P. acknowledges the support from NSF grant DMR-1005932.

¹ G. Jackeli and G. Khaliullin, Phys. Rev. Lett. **102**, 017205 (2009).

² J. Chaloupka, G. Jackeli, and G. Khaliullin, Phys. Rev. Lett. **105**, 027204 (2010).

³ G.-W. Chern and N. B. Perkins, Phys. Rev. B **80**, 180409(R) (2009).

⁴ *Spin liquids in frustrated magnets*, Leon Balents, Nature **464**, 199 (2010).

- ⁵ A. Kitaev, *Ann. Phys.* **321**, 2 (2006).
- ⁶ Y. Singh and P. Gegenwart, *Phys. Rev. B* **82**, 064412 (2010).
- ⁷ X. Liu, T. Berlijn, W.-G. Yin, W. Ku, A. Tsvelik, Young-June Kim, H. Gretarsson, Yogesh Singh, P. Gegenwart, and J. P. Hill, *Phys. Rev. B* **83**, 220403 (2011).
- ⁸ Yogesh Singh, S. Manni, J. Reuther, T. Berlijn, R. Thomale, W. Ku, S. Trebst, P. Gegenwart *Phys. Rev. Lett.* **108**, 127203 (2012).
- ⁹ S. K. Choi, R. Coldea, A. N. Kolmogorov, T. Lancaster, I. I. Mazin, S. J. Blundell, P. G. Radaelli, Yogesh Singh, P. Gegenwart, K. R. Choi, S.-W. Cheong, P. J. Baker, C. Stock, and J. Taylor, *Phys. Rev. Lett.* **108**, 127204 (2012).
- ¹⁰ H.-C. Jiang, Z.-C. Gu, X.-L. Qi, and Simon Trebst *Phys. Rev. B* **83**, 245104 (2011).
- ¹¹ J. Reuther, R. Thomale, and S. Trebst, *Phys. Rev. B* **84**, 100406 (2011).
- ¹² F. Trouselet, G. Khaliullin, P. Horsch, *Phys. Rev. B* **84**, 054409 (2011).
- ¹³ Timo Hyart, Anthony R. Wright, Giniyat Khaliullin, and Bernd Rosenow *Phys. Rev. B* **85**, 140510 (2012).
- ¹⁴ Crag Price and Natalia B. Perkins, *Phys. Rev. Lett.* **109**, 187201 (2012).
- ¹⁵ I. I. Mazin, H. O. Jeschke, R. Valenti, D. I. Khomskii, *Phys. Rev. Lett.* **109**, 197201 (2012).
- ¹⁶ R. Schaffer, S. Bhattacharjee, Y. B. Kim, *Phys. Rev. B* **86**, 224417 (2012).
- ¹⁷ I. Kimchi and Y.Z. You, *Phys. Rev. B* **84**, 180407(R) (2011).
- ¹⁸ SubhroBhattacharjee, Sung-Sik Lee, Yong Baek Kim, *New J. Phys.* **14**, 073015 (2012).
- ¹⁹ J. Chaloupka, G. Jackeli, G. Khaliullin, *PRL* **110**, 097204 (2013).
- ²⁰ N. D. Mermin and H. Wagner, *Phys. Rev. Lett.* **17**, 1133 (1966).
- ²¹ J.V. José, L.P. Kadanoff, S. Kirkpatrick S and D. R. Nelson, *Phys. Rev. B* **16**, 1217 (1977).
- ²² G. Ortiz, E. Cobanera, Z. Nussinov, *Nuclear Physics B* **854**, 780 (2012).
- ²³ G. Baskaran, Diptiman Sen, and R. Shankar, *Phys. Rev. B* **78**, 115116 (2008).
- ²⁴ R. J. Baxter, *J. Math. Phys.* **11**, 784 (1970); P. W. Kasteleyn, *J. Math. Phys.* **4**, 287 (1963).
- ²⁵ It is interesting to note that emergent U(1) symmetry of discrete order parameter has been recently widely discussed in the literature.^{26–30} For example, an emergent U(1) symmetry is a major prediction of the theory of the classical 120° orbital model on the simple cubic lattice,²⁸ it appears from the non-trivial decoupling of the Heisenberg antiferromagnet on interpenetrating honeycomb and triangular lattices,³⁰ and is also present in the theory of deconfined critical points for dimer order parameters in the 2D Heisenberg model with four-spin interactions.²⁹
- ²⁶ S. V. Isakov and R. Moessner, *Phys. Rev. B* **68**, 104409 (2003).
- ²⁷ G.-W. Chern, O. Tchernyshyov, *Phil. Trans. Roy. Soc. A* **370**, 5718 (2012)
- ²⁸ Sandro Wenzel, Andreas M. Laeuchli, *Journal of Statistical Mechanics: Theory and Experiment* 2011, P09010 (2011).
- ²⁹ Anders W. Sandvik, *Phys. Rev. Lett.* **98**, 227202 (2007).
- ³⁰ Peter P. Orth, Premala Chandra, Piers Coleman, Jrg Schmalian *Phys. Rev. Lett.* **109**, 237205 (2012)
- ³¹ M.S.S. Challa and D.P. Landau, *Phys. Rev. B* **33**, 437 (1986).
- ³² J. M. Kosterlitz, *J. Phys. C* **7**, 1046 (1974).